

# Delphinium

## 30ct Plug

Container:

Trade 1 Gallon Container

Finish:

6-8 weeks

When To Plant:

Early to late spring

## 72ct Plug

Container:

Trade 1 Gallon Container

Finish:

8-10 weeks

When To Plant:

Early to late spring

### Growing Temperature:

65-75° F days

50-55° F nights

### EC Level:

1.5-2.0 pour through method

### Vernalization:

Not required.

### Pests and Diseases:

Aphids, borers, leaf miner, mites (cyclamen, brood, two-spotted spider), lily aphid, green peach aphid (Myzus) in greenhouse, beetles, cutworm, sowbugs, slugs, nematodes, aster leafhopper, thrips.

Impatiens necrotic spot virus, viruses, aster yellows, Verticillium wilt, bacterial blight (Pseudomonas), stem canker (Fusarium), damping off, smut, crown rots, Botrytis blight, black leg (Erwinia), crown gall, leaf spots, powdery mildew, rusts, chlorosis (low temperature, wet soil).

Watch for powdery mildew, botrytis and erwinia soft rot when high humidity and temperatures occur. Avoid stagnant air conditions, especially when flower buds are forming. A general fungicide treatment is advisable if these environmental conditions persist.

### Lighting:

For best results grow under photoperiods of 12 hours or longer.

### Holding Temperature:

50-55° F

### Soil pH:

5.8-6.2

### Fertility:

A balanced fertilizer using calcium and potassium nitrate is preferred. Once flower initiation occurs, use less nitrogen and more potassium and phosphorus. Feed 75-100 ppm N at each watering.

For Controlled Release or Slow Release Fertilizer, see your preferred supplier for recommended rates for incorporation or top dressing, as it varies by fertilizer.

### Potting and Timing:

Plant in early to late spring.

### Moisture:

Water well when first planted, then keep damp to moist, but not soggy, until initial bud breaks. Do not let the soil dry out completely. Avoid overhead watering to prevent diseases. Avoid overwatering and wet foliage after flower bud initiation occurs.

### Planting Level:

Place crown at or just below soil surface.

### PGRs/Pinching:

Apply Bonzi at 30 ppm or Sumagic at 5 ppm as a spray when flower stems begin to grow beyond the foliage.

### Grower Tips:

Good air circulation is important to reduce the chance of disease problems.



P.O. Box 137 • 1992 96th Ave. Zeeland, MI. 49464  
Phone: 888-925-8377 • Fax: 800-752-1879  
Email: [sales@waltersgardens.com](mailto:sales@waltersgardens.com)  
[www.WaltersGardens.com](http://www.WaltersGardens.com) • [www.PerennialResource.com](http://www.PerennialResource.com)  
[www.ProvenWinners.com](http://www.ProvenWinners.com)